

ARMEN VARTANY, M. D.
Certified, American Board of Plastic Surgery
116 South Buena Vista Street
Burbank, California 91505
(818) 500-0823

INSTRUCTIONS FOR ABDOMINOPLASTY with LIPOSUCTION

BEFORE SURGERY

Do not take aspirin, Excedrin, Ibuprofen, Motrin, Advil, Aleve or any other non-steroidal anti-inflammatories, for two weeks prior to surgery and for one week after surgery. These medications interfere with normal blood clotting. You may take Tylenol, if needed, for minor aches.

Discontinue the use of Birth Control pills for two weeks prior to surgery. You may resume taking your birth control one week after surgery.

Self-Quarantine after the COVID-19 test has been administered during your pre-op.

For a period of two days prior to surgery, use antibacterial soap (Dial, Safeguard, Phase III, etc.), for bathing.

Telephone the office if you notice any signs of fever, infection, pustules, or a cold.

You may brush your teeth the morning of surgery; but **have nothing to eat or drink after midnight the night before surgery.** If you take any medications for blood pressure or your heart, please take them with a sip of water as scheduled.

On the day of surgery, wear a loose fitting, button-up, blouse, or shirt type dress, which has been freshly laundered. Avoid any article of clothing that requires being put on over the head. Do not wear pantyhose, boots, and girdles. Loose, lounge pants and simple slip-on shoes or slippers.

Do not wear any make-up, nail polish or jewelry to the surgery center.

Wear a face mask or protective covering to the surgery center.

If you wear contact lenses, be sure to bring a case and solution as your contact lenses will need to be removed prior to surgery. You will be able to put your contact lenses back in approximately one day after surgery.

Please park on the second floor of our parking structure. This is the level that you will be discharged.

When you arrive for surgery, you will advise the facility who is to be called to take you home. **Your companion may not enter the office or wait in the office during your surgery (unless you are a minor).** Once you have been transitioned to the recovery room, your designated ride will be contacted. Your ride will need to arrive within 30 minutes for instructions and discharge.

You **must** have someone spend the first night with you.

If you have any questions, prior to surgery, please call the office any time.

AFTER SURGERY:

Return home and go directly to bed, with only assisted bathroom privileges for the first 24 hours.

Take medications as directed**. Remember that NO alcohol is permitted for one week to ten days, depending on when you are allowed to cease taking medications.

Upon arrival to your home, you may have small quantities of liquids as desired and/or tolerated. It is advised that you eat a very small amount of pudding or yogurt to coat the stomach prior to taking pain medication. **If you are experiencing any nausea, avoid taking the anti-biotic and/or pain medication until the nausea subsides or until the following day when you may progress to a soft diet. After that, you may eat as desired and/or tolerated. Hot and/or spicy foods may cause additional swelling.

You will have to empty your surgical drain(s) twice daily and write down the amount of fluid removed from each drain and each time. Your drains will be removed in the office when the output goes down, which is usually 7 to 10 days after your surgery.

Do not shower while the drains are in place. You may have a sponge bath on the second day after surgery, but under no circumstances, should the bandages or drains become wet. You will be able to take a shower after your drains have been removed.

Your abdominal girdle should be worn for six weeks after the operation. Do not remove your girdle unnecessarily. After your drain(s) has been removed, you may remove the girdle to wash, but then place back on. The girdle will help with your new shape, prevent accumulation of fluid and support the healing tissues.

Do not attempt to sleep on your stomach for four to six weeks after surgery.

Avoid getting overheated, and do not sit out in the sun for at least two weeks. Additionally, avoid any exercise and/or strenuous activity that could potentially elevate your heart rate or blood pressure for four weeks after surgery.

Do not drive a car until you are given permission.

If there are any signs of persistent nausea, bleeding, rash, etc., you are to report this to the office.

Should you have a question or concern during office hours:

Monday – Friday 7:30am – 6pm (Closed 12pm - 1pm for lunch)

Saturday 7:30am – 4pm (Closed 12pm – 1pm for lunch)

Please call the office at (818) 500-0823.